

1993

## High-Velocity Formaldehyde Absorption Within 30" of the Galactic Nucleus

T. Pauls

K. J. Johnston

T. L. Wilson

Jonathan Marr

*Haverford College, jmarr@haverford.edu*

Follow this and additional works at: [http://scholarship.haverford.edu/physics\\_facpubs](http://scholarship.haverford.edu/physics_facpubs)

---

### Repository Citation

"High-Velocity Formaldehyde Absorption Within 30" of the Galactic Nucleus" by T. Pauls, K. J. Johnston, T. L. Wilson, J. M. Marr, and A. Rudolph 1993, *Astrophys. J.*, 403, L13

This Journal Article is brought to you for free and open access by the Physics at Haverford Scholarship. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Haverford Scholarship. For more information, please contact [nmedeiro@haverford.edu](mailto:nmedeiro@haverford.edu).

## HIGH-VELOCITY FORMALDEHYDE ABSORPTION WITHIN 30" OF THE GALACTIC NUCLEUS

T. PAULS,<sup>1</sup> K. J. JOHNSTON,<sup>1</sup> T. L. WILSON,<sup>2</sup> J. M. MARR,<sup>3</sup> AND A. RUDOLPH<sup>4</sup>

Received 1992 February 24; accepted 1992 November 5

### ABSTRACT

We report VLA observations of the  $(1_{10}-1_{11})$  line of  $\text{H}_2\text{CO}$  in absorption within 30" of the Galactic center point source Sagittarius A\* (Sgr A\*) with a resolution of  $20''.6 \times 9''.5$  at P.A. =  $4^\circ$ , and an rms noise of  $3.5 \text{ mJy beam}^{-1}$ . The absorption peaks within 15" of Sgr A\*, and extends in velocity from about  $+135 \text{ km s}^{-1}$  to at least  $-210 \text{ km s}^{-1}$  (with some interruptions). We performed a number of baseline subtractions using different velocity intervals to define the continuum level. In all cases the absorption is definitely present. At positions offset  $\pm 1'$  in latitude from Sgr A\* there is no evidence for absorption at high velocities. Thus, the high-velocity gas may be close to, but is not coincident with, Sgr A\*. The gas with velocities between  $-157$  and  $-205 \text{ km s}^{-1}$  is located east and northeast of the continuum peak, and gas at about  $+135 \text{ km s}^{-1}$  is also located to the east. The position of the absorption relative to the continuum leads us to believe that the negative velocity molecular gas is behind Sgr A\*. Because the gas is localized near Sgr A\*, we do not think that there is any interaction with Sgr A East.

Taken together with observations of  $\text{HCO}^+$ ,  $\text{HCN}$ , and  $[\text{O I}]$ , there are now several lines of evidence which indicate that the central cavity at the Galactic center contains a mixture of neutral and ionized gas.

*Subject headings:* Galaxy: center — galaxies: nuclei — ISM: molecules

### 1. INTRODUCTION

The region within a radius of 30" of the Galactic nucleus contains neutral and ionized gas clouds, stars and star clusters, magnetic fields, and a radio point source, Sgr A\*, believed to be the dynamical center of the Galaxy (cf. Genzel & Townes 1987, Morris 1989, and references therein). At larger radii,  $r \sim 2'-3'$ , the radio radiation from the Galactic center shows an extended region of nonthermal emission called Sgr A East, and a somewhat more compact source of thermal emission, Sgr A West. The region around Sgr A\*, described above, is associated with Sgr A West. We follow Pedlar et al. (1989) in placing Sgr A East, thought to be a supernova remnant, behind Sgr A West. Spectroscopic studies at radio and infrared wavelengths of the gas kinematics reveal large-scale systematic motions with velocities, both positive and negative, up to several hundred  $\text{km s}^{-1}$ . The ionized gas, studied via continuum radiation and emission lines of  $\text{H}^+$  (Roberts et al. 1991) and  $\text{Ne}^+$  (Lacy, Achterman, & Serabyn 1991) at  $1''-2''$  resolution, is concentrated in a spiral-like, filamentary structure centered on Sgr A\* (Lo & Claussen 1983). Lacy et al. have modeled the kinematics of the ionized gas using approximately circular orbits with a nearly Keplerian rotation curve which requires a central point mass of  $2 \times 10^6 M_\odot$ . It is generally thought that within  $\leq 1.5 \text{ pc}$  (about 40" if  $R_0 = 8.5 \text{ kpc}$ ) there is a cavity filled primarily with ionized gas. However, recent observations of the  $[\text{O I}] 63 \mu\text{m}$  line show that neutral atomic gas is present in the central cavity (Jackson et al. 1993). The molecular gas is thought to lie primarily outside the cavity and appears to delineate a clumpy ring or disk with an inner radius of  $\sim 1.5 \text{ pc}$  and a rotational velocity of about  $100 \text{ km s}^{-1}$  (Güsten et al. 1987).

In spite of all these data, until recently, the question of outflow versus inflow of molecular gas toward Sgr A\* was still open. Zylka, Mezger, & Wink (1990) and Ho et al. (1991) have argued from 10" and 12" maps that molecular gas *must* be flowing toward Sgr A\*. However, large negative velocity gas is found in absorption against the continuum. The most notable example is the absorption of  $\text{H}_2\text{CO}$  between  $-160$  and  $-200 \text{ km s}^{-1}$  at 4.8 GHz observed by Güsten & Downes (1981) with 3' resolution. They concluded that this gas may have been ejected from the nucleus.

In this *Letter* we present new VLA observations of  $\text{H}_2\text{CO}$  absorption made with a resolution of  $20''.6 \times 9''.5$ . Our data give the location of the  $\text{H}_2\text{CO}$  absorbing gas relative to the radio continuum from the ionized gas.

### 2. OBSERVATIONAL PARAMETERS AND DATA REDUCTION

The Galactic center was observed on 1988 August 19 with the D configuration of the NRAO<sup>5</sup> VLA for a 7 hr period at the frequency of the  $\text{H}_2\text{CO}(1_{10}-1_{11})$  transition, 4.82966 GHz. The spectrometer bandwidth was 6.25 MHz, divided over 128 channels, giving a velocity resolution of  $3 \text{ km s}^{-1}$  (no Hanning smoothing), and a velocity coverage from  $-230$  to  $+150 \text{ km s}^{-1}$ . Observations of 3C 286 (7.46 Jy at 4.8 GHz) were used to calibrate the spectrometer bandpass and set the flux density scale, and NRAO 530 was used for phase calibration. All data reduction and analysis was done using the Astronomical Image Processing System (AIPS). The continuum level was established by using the emission between  $-70$  and  $-100 \text{ km s}^{-1}$  and between  $+102$  and  $+121 \text{ km s}^{-1}$ . After calibration, and bandpass correction, the continuum emission was subtracted from the  $u-v$  data base using the task UVBAS (van Langevelde & Cotton 1990). The resulting line-minus-continuum  $u-v$  data were edited, mapped onto a  $256 \times 256$  grid with a  $3'' \times 3''$  pixel size using uniform weighting, and deconvolved,

<sup>1</sup> Remote Sensing Division, Naval Research Laboratory, Washington, DC 20375.

<sup>2</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-5300, Bonn 1, Germany.

<sup>3</sup> Astronomy Department, Haverford College, Haverford, PA 19041.

<sup>4</sup> Department of Astronomy, University of Maryland, College Park, MD 20742.

<sup>5</sup> The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under cooperative agreement with the National Science Foundation.

yielding a synthesized beam of  $20''.6 \times 9''.5$  at P.A. =  $4^\circ$ , and a final rms noise on the individual channel maps of about  $3.5 \text{ mJy beam}^{-1}$ .

### 3. RESULTS AND DISCUSSION

#### 3.1. Comparison with the Single-Dish Spectrum

Figure 1 shows a comparison between the single-dish  $\text{H}_2\text{CO}$  absorption spectrum made by Güsten & Downes with  $3'$  resolution and our VLA data averaged over  $15'' \times 15''$ , both centered on Sgr A\*. In this Letter we will discuss high-velocity absorption confined within  $\pm 30''$  of Sgr A\*. Inspection of Figure 1 shows that there is absorption from  $+135 \text{ km s}^{-1}$  to beyond  $-210 \text{ km s}^{-1}$ , with some interruptions, and is nearly continuous from  $-120 \text{ km s}^{-1}$  to about  $-210 \text{ km s}^{-1}$ , where we have cut off the plot because of the increased noise near the edge of the spectrometer bandpass. The  $\text{H}_2\text{CO}$  spectrum of Güsten & Downes shows absorption between  $-70$  and  $-100 \text{ km s}^{-1}$  which is not present in our spectrum. We used data in this velocity range and at velocities between  $+102$  and  $+121 \text{ km s}^{-1}$  to determine the continuum level in our data, and, thus the absolute value of the absorption line we observe may be underestimated. In order to test the sensitivity of the positions of the absorption features to the choice of continuum channels, we have reanalyzed our data using only channels between  $+102$  and  $+139 \text{ km s}^{-1}$  to define the continuum level. These tests show that the absorption peaks shift by 1–3 pixels, and we conclude that the positions of the absorption features should be accurate to about  $10''$ . In addition, the absorption may extend over more than the  $\sim 350 \text{ km s}^{-1}$  presented here, as is suggested by the line shape seen in Figure 1.

From Figure 2 (Plate L1), a longitude-velocity diagram of  $\text{H}_2\text{CO}$  absorption made at the latitude of Sgr A\*, we see that there is extensive absorption within  $\pm 30''$  of Sgr A\*. Similar longitude-velocity diagrams made at latitudes offset  $\pm 1'$  from the latitude of Sgr A\* show no high-velocity absorption. However, assuming all the continuum is behind the molecular clouds, it is possible that formaldehyde is present at these velocities but is not detected due to the lower continuum levels at




FIG. 1.—Composite  $\text{H}_2\text{CO}$  absorption spectrum showing the single-dish data from Güsten & Downes (1981) in a  $3'$  beam (dotted line) divided by 10 and our data averaged over a  $15'' \times 15''$  region, both centered on Sgr A\*. Note that there are many points at which the spectra do not agree. This is especially true in the  $-50$  to  $-130 \text{ km s}^{-1}$  range.

the offset positions. Figures 1 and 2 suggest that the  $\text{H}_2\text{CO}$  clouds have a fairly clumpy distribution, with several narrow peaks of increased absorption superposed on broader, weak absorption. Some of the peaks are associated with well-known features seen along the line of sight to the Galactic center such as the “3 kpc arm” at  $-50 \text{ km s}^{-1}$ , and the “expanding molecular ring” at  $-135 \text{ km s}^{-1}$ , which is located about 200 pc from the Galactic center. The absorption which peaks at around  $+90 \text{ km s}^{-1}$  is seen in OH (Sandqvist et al. 1987) and HCN (Güsten et al. 1987) at the position of the circumnuclear ring. Although Whiteoak, Gardner, & Pankonin (1983) have published maps of Sgr A in the 4.8 GHz  $\text{H}_2\text{CO}$  line, features at  $-170$  to  $-200 \text{ km s}^{-1}$  were not detected because of the lower sensitivity of those observations. We identify the weak, broad absorption at negative velocities with the “high-velocity gas” seen by Güsten & Downes (1981).

#### 3.2. The Location of the Absorbing Gas

In Figure 3 we plot absorption maps of the negative velocity channels from  $-167$  to  $-213 \text{ km s}^{-1}$ . Figures 2 and 3 show that the peaks of the high-velocity gas lie within about  $\pm 15''$  of Sgr A\*. The blueshifted clouds are primarily located east and northeast of the nucleus, although there is gas to the west at  $-172 \text{ km s}^{-1}$  and weaker absorption about  $25''$  southwest of Sgr A\* between  $-185$  and  $-192 \text{ km s}^{-1}$ . The redshifted gas at  $+135 \text{ km s}^{-1}$  also peaks to the east. *In no case does the maximum absorption appear to be coincident with Sgr A\**. We note that most of the peaks in Figure 3 are unresolved in our  $20''.6 \times 9''.5$  beam.

In order to investigate the relationship between the neutral gas, traced by the  $\text{H}_2\text{CO}$ , and the ionized gas in the nucleus, we have made the composite maps in Figure 4 (Plate L2). Here the  $\text{H}_2\text{CO}$  absorption is displayed as a gray-scale image superposed on contours of our 4.8 GHz continuum, which includes emission from the ionized filaments smoothed by our beam. Within the limits imposed by the spatial resolution of the present observations, we feel Figure 4 suggests that the neutral gas may lie between the ionized filaments, but this suggestion needs to be verified by observations with higher spatial resolution. Assuming that all the continuum is behind the  $\text{H}_2\text{CO}$  clouds, we derive apparent optical depths,  $\tau \sim 0.01$ – $0.02$ , which implies  $A_v \sim 1$ – $2$  mag. Thus, the actual optical depths may be larger. If  $T_{\text{ex}}$  is constant, then our absorption maps also trace column density.

Figures 3 and 4 show that the maximum  $\text{H}_2\text{CO}$  absorption occurs at projected distances as close as  $15''$ – $20''$  from Sgr A\*. We feel that the small projected distance and the high radial velocity strongly suggest that the neutral gas traced by  $\text{H}_2\text{CO}$  is near the Galactic nucleus, well within the 10 pc lower limit set by Güsten & Downes (1981). In fact, it may well be that the projected and true distances are very nearly the same. For example, if the gas originates in the circumnuclear disk with a rotational velocity of  $\sim 110 \text{ km s}^{-1}$  at about 1.5 pc from the center and falls in the gravitational field of a  $4 \times 10^6 M_\odot$  point mass, it will reach a rotational velocity of  $\sim 200 \text{ km s}^{-1}$  at a distance of 0.6 pc from the nucleus. Since the observed velocity is only the radial component of the total velocity, this gas is probably closer than 0.6 pc from Sgr A\*.

However, Figure 3 shows that none of the  $\text{H}_2\text{CO}$  absorption maxima peak at the position of Sgr A\*. The maximum in the  $\text{HCO}^+$  absorption at large negative velocities reported by Marr et al. (1992) is offset to the southwest and west of Sgr A\*, and an H I spectrum with  $3''$  resolution toward Sgr A\* shows




FIG. 2.—Longitude-velocity diagram of  $\text{H}_2\text{CO}$  absorption at the latitude of Sgr A\* ( $b = -2.75$ ). Contour levels are  $-10$ ,  $-20$ ,  $-30$ , and  $-40 \text{ mJy beam}^{-1}$ . Note the weak, high-velocity absorption appears to be confined within about  $\pm 30''$  of Sgr A\*. There appear to be two sets of extended, high-velocity absorption features located on either side of  $\Delta l = 0$ .

PAULS et al. (see 403, L14)




FIG. 4.—(a) A composite figure showing the  $\text{H}_2\text{CO}$  absorption at  $-188.5 \text{ km s}^{-1}$  superposed on the 4.8 GHz continuum from Sgr A. The position of Sgr A\* is marked by a cross. The contour levels for the continuum are 5, 10, 20, 40, 60, and 80% of the peak flux of  $4.07 \text{ Jy beam}^{-1}$ . The gray scale at the top of each plot shows the absorption in  $\text{mJy beam}^{-1}$ . (b) The  $\text{H}_2\text{CO}$  absorption at  $+132.8 \text{ km s}^{-1}$ .

PAULS et al. (see 403, L14)






FIG. 3.—Individual channel maps of H<sub>2</sub>CO absorption for velocities from  $-167$  to  $-213$  km s<sup>-1</sup>. Note that the absorption is located primarily east and northeast of Sgr A\* (denoted by the cross). The synthesized beam is shown as an ellipse in the channel at  $-167$  km s<sup>-1</sup>. Contour levels are  $-20$ ,  $-30$ , and  $-40$  mJy beam<sup>-1</sup>.

no absorption at velocities beyond  $-150$  km s<sup>-1</sup> (Liszt, Burton, & van der Hulst 1985). All these data suggest that high-velocity neutral gas lies close to the position of Sgr A\* but is not coincident with it.

If the nonthermal source Sgr A East lies behind Sgr A West (Pedlar et al. 1989), and Sgr A\* is close to Sgr A West, then the neutral gas we observe may be between Sgr A East and Sgr A West. Additionally, if Sgr A East is a supernova remnant (SNR), then the high negative radial velocities might be caused by molecular gas in front of Sgr A East, pushed in our direction by the SNR shell. In this case there might be little or no connection with Sgr A\*. However, our  $l$ - $v$  diagrams at latitude offsets of  $\pm 1'$  from Sgr A\* show no evidence of high negative velocity gas. We feel, therefore, that the blueshifts may arise near Sgr A\*.

#### 4. CONCLUDING REMARKS

There are now several independent lines of evidence which indicated that neutral gas may be mixed with ionized gas

within 0.5 pc of the Galactic nucleus. The neutral gas has now been detected in H<sub>2</sub>CO (this paper), HCO<sup>+</sup> (Marr et al. 1992), and HCN and [O I] (Jackson et al. 1993); and there is now strong evidence that massive star formation has occurred in the central parsec within the last few million years (Krabbe et al. 1991), producing a cluster of young, hot stars. In addition, recent infrared polarimetry observations at  $12.4 \mu\text{m}$  demand the presence of dust grains in this region as well (Aitken et al. 1991). If the dust provides the shielding so that photo-dissociation regions form (Tielens & Hollenbach 1985), our results support the suggestion that the northern and eastern arms of the ionized filaments may be bright ionized rims on the surface of the neutral gas clouds (Jackson et al. 1993).

Our data are consistent with a mass within 0.4 pc of  $3 \times 10^6 M_{\odot}$ , derived from other tracers, but do not in any way place a limit on the central mass of Sgr A. These results show that the main objections to infall are no longer valid: the *negative velocity* gas seems to be behind Sgr A\*. However, any infall model must satisfy rather complex kinematic conditions.

#### REFERENCES

- Aitken, D. K., Gezari, D., Smith, C. H., McCaughrean, M., & Roche, P. F. 1991, ApJ, 380, 419  
 Genzel, R., & Townes, C. H. 1987, ARA&A, 25, 377  
 Güsten, R., & Downes, D. 1981, A&A, 99, 27  
 Güsten, R., Genzel, R., Wright, M. C. H., Jaffe, D. T., Stutzki, J., & Harris, A. I. 1987, ApJ, 318, 124  
 Ho, P. T. P., Ho, L. C., Szczepanski, J. C., Jackson, J. M., Armstrong, J. T., & Barrett, A. H. 1991, Nature, 350, 309  
 Jackson, J. M., Geis, N., Genzel, R., Harris, A. I., Madden, S. C., Poglitsch, A., Stacey, G. J., & Townes, C. H. 1993, ApJ, 402, 173  
 Krabbe, A., Genzel, R., Drapatz, S., & Rotaciuc, V. 1991, ApJ, 382, L19  
 Lacy, J. H., Achtermann, J. M., & Serabyn, E. 1991, ApJ, 380, L71  
 Liszt, H. S., Burton, W. B., & van der Hulst, J. M. 1985, A&A, 142, 245  
 Lo, K.-Y., & Claussen, M. J. 1983, Nature, 306, 647  
 Marr, J. M., Rudolph, A., Pauls, T. A., Wright, M. C. H., Backer, D. C., & Tucker, L. M. 1992, ApJ, 400, L29

L16

PAULS ET AL.

- Morris, M., ed. 1989, *The Center of the Galaxy* (Dordrecht: Kluwer)
- Pedlar, A., Anantharamaiah, K. R., Ekers, R. D., Goss, W. M., van Gorkom, J. H., Schwartz, U. J., & Zhao, J.-H. 1989, *ApJ*, 342, 769
- Roberts, D. A., Goss, W. M., van Gorkom, J. H., & Leahy, J. P. 1991, *ApJ*, 366, L15
- Sandqvist, A., Karlsson, R., Whiteoak, J. B. & Gardner F. F. 1987, in *The Galactic Center, Proc. of a Symposium Honoring C. H. Townes*, ed. D. Backer (New York: AIP), 95
- Serabyn, E., Güsten, R., Walmsley, C. M., Wink, J. E., & Zylka, R. 1986, *A&A*, 169, 85
- Tielens, A. G. G. M., & Hollenbach, D. 1985, *ApJ*, 291, 722
- van Langevelde, H. J., & Cotton, W. D. 1990, *A&A*, 239, L5
- Whiteoak, J. B., Gardner, F. F., & Pankonin, V. 1983, *MNRAS*, 202, 11P
- Zylka, R., Mezger, P. G., & Wink, J. E. 1990, *A&A*, 234, 133